

QUICK REFERENCE VACCINATION SHEET

Vaccinations are to be given as **single doses**, never together in the same appointment, and always spread out 4 weeks from any other medical procedure or other vaccine. A sick puppy should NEVER be vaccinated.

Age	Vaccine	Frequency	Booster
12-14 weeks	Parvovirus	ONCE	Never
16-20 weeks (or 4 weeks after parvo shot)	Distemper*	ONCE	Never
12 months	Rabies	ONCE	Never**

*Distemper is KNOWN to cause serious and lifelong reactions in Tollers. This vaccine should only be administered if there is a SERIOUS likelihood of your dog coming in contact with diseased animals.

**If you live in the USA or elsewhere and a rabies vaccination is required by law, ensure you are asking for the 3 year vaccine to extend the amount of time necessary between shots.

Things you can do to alleviate some of the stress vaccination places on your puppy:

- Give Vitamin C to the puppy for several days before and after his shot
- Give the Homeopathic Thuja in 30 cc following the shot
- Check your local dog health food stores for Pet Innoculant by Wysong- give orally

No other vaccines or boosters required **ever**.

[DO NOT ALLOW YOUR VET TO TALK YOU INTO ADDITIONAL VACCINATIONS WITHOUT FIRST DISCUSSING WITH YOUR BREEDER. ADMINISTERING VACCINES NOT AUTHORIZED BY YOUR BREEDER WILL VOID YOUR HEALTH WARRANTY.]

Annual titre testing can prove persistent immunity to the diseases for which your dog was vaccinated. Contact your breeder IMMEDIATELY if you are being asked for proof of vaccination to attend a class, kennel, daycare, boarding facility, groomer, etc.

